

Mitochondrial DNA Part B Resources

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

The complete chloroplast genome of dalmatian pyrethrum (*Tanacetum cinerariifolium* (Trevir.) Sch. Bip. (Asteraceae)), the source of the natural insecticide pyrethrin

Ante Turudić, Zlatko Liber, Martina Grdiša, Jernej Jakše, Filip Varga & Zlatko Šatović

To cite this article: Ante Turudić, Zlatko Liber, Martina Grdiša, Jernej Jakše, Filip Varga & Zlatko Šatović (2022) The complete chloroplast genome of dalmatian pyrethrum (*Tanacetum cinerariifolium* (Trevir.) Sch. Bip. (Asteraceae)), the source of the natural insecticide pyrethrin, *Mitochondrial DNA Part B*, 7:5, 775-777, DOI: [10.1080/23802359.2022.2070439](https://doi.org/10.1080/23802359.2022.2070439)

To link to this article: <https://doi.org/10.1080/23802359.2022.2070439>



© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 05 May 2022.



Submit your article to this journal [↗](#)



Article views: 303



View related articles [↗](#)



View Crossmark data [↗](#)

The complete chloroplast genome of dalmatian pyrethrum (*Tanacetum cinerariifolium* (Trevir.) Sch. Bip. (Asteraceae)), the source of the natural insecticide pyrethrin

Ante Turudić^{a,b}, Zlatko Liber^{b,c}, Martina Grdiša^{a,b}, Jernej Jakše^d, Filip Varga^{a,b} and Zlatko Šatović^{a,b}

^aCentre of Excellence for Biodiversity and Molecular Plant Breeding (CoE CroP-BioDiv), Zagreb, Croatia; ^bFaculty of Agriculture, University of Zagreb, Zagreb, Croatia; ^cFaculty of Science, University of Zagreb, Zagreb, Croatia; ^dBiotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia

ABSTRACT

Tanacetum cinerariifolium is an endemic species of the eastern Adriatic coast that synthesizes the natural insecticide pyrethrin. We have characterized the complete chloroplast genome of the species and analyzed its phylogeny within the Asteraceae family. The complete chloroplast genome of *T. cinerariifolium* has a size of 150,136 bp, including a large single-copy (LSC) region of 82,717 bp, a small single-copy (SSC) region of 18,411 bp, and a pair of inverted repeats (IRs) of 24,504 bp. The chloroplast genome of *T. cinerariifolium* encodes 108 genes, including 77 protein-coding genes (PCGs), 27 tRNA genes, and 4 rRNA genes. Phylogenetic analyses based on the complete chloroplast genomes placed *T. cinerariifolium* in a sister position to species of the genera *Artemisia* and *Chrysanthemum*.

ARTICLE HISTORY

Received 15 October 2021

Accepted 15 April 2022

KEYWORDS

Tanacetum cinerariifolium;
chloroplast genome; phylogeny

Dalmatian pyrethrum (*Tanacetum cinerariifolium* (Trevir.) Sch. Bip. 1844) is a perennial, thermophilous plant species in the family Asteraceae. Autochthonous populations of *T. cinerariifolium* are distributed along the coast of the eastern Adriatic Sea, mainly in Croatia, Montenegro, and Albania (Nikolić 2020). The plant species synthesizes the secondary metabolite pyrethrin, which is known worldwide for its effective insecticidal and insect repellent activity, with little to no negative impact on humans and the environment (Casida and Quistad 1995). In this study, we assembled the complete chloroplast sequence of *T. cinerariifolium* and subjected it to phylogenetic analysis.

The plant of *T. cinerariifolium* was collected near Vrbnik, the island of Krk, Croatia (45°04'33" N, 14°40'21" E; 45 m a.s.l.). The voucher specimen was deposited in the Herbarium ZAGR of the University of Zagreb, Faculty of Agriculture, Zagreb, Croatia under Herbarium ID 47,683 (available at: <http://herbarium.agr.hr>, contact Sandro Bogdanović, sbogdanovic@agr.hr).

Total cellular DNA was isolated from 100 mg of fresh leaf tissue using the DNeasy Plant Mini Kit (Qiagen GmbH, Hilden, Germany). Sequencing was performed using Illumina NovaSeq600 (Illumina, Inc., San Diego, CA). The base calls (BCL) binary was converted to FASTQ using the Illumina package bcl2fastq.

Whole genome sequencing yielded a total of 796,130,142 reads and the total base reads were 120.2 Gbp.

The chloroplast genome was assembled using the Fast-Plast version 1.2.8 pipeline (McKain and Wilson 2018). In Fast-

Plast, reference genomes were set to order Asterales. Annotation was performed using the online program GeSeq (Tillich et al. 2017).

The genome was 150,136 bp long and consisted of four distinct regions, including a large single-copy region (LSC; 82,717 bp), a small single-copy region (SSC; 18,411 bp), and a pair of inverted repeat regions (IRs; 24,504 bp each). The genome contained 108 unique genes, including 77 protein-coding genes (PCGs), 27 tRNA genes, and 4 rRNA genes. The total GC content was 37.4%, while the corresponding value for the LSC, SSC, and IR regions was 35.5%, 30.8%, and 43.1%, respectively.

The chloroplast genome sequences of 14 species of Asteraceae family were downloaded from the NCBI Nucleotide database and used for phylogenetic analysis, including 13 species from the subfamily Asteroideae (10 from tribe Anthemideae Cass. and three from tribe Astereae Cass.) and the outgroup from the subfamily Cichorioideae: *Anaphalis sinica* Hance (NC_034648), *Archibaccharis asperifolia* (Benth.) S.F.Blake (NC_034848), *Artemisia argyi* H.Lev. & Vaniot (NC_030785), *Artemisia capillaris* Thunb. (NC_031400), *Artemisia frigida* Willd. (NC_020607), *Artemisia gmelinii* Weber ex Stechm. (NC_031399), *Artemisia montana* (Nakai) Pamp. (NC_025910), *Aster altaicus* Willd. (NC_034996), *Chrysanthemum boreale* (Makino) Makino (NC_037388), *Chrysanthemum indicum* L. (NC_020320), *Chrysanthemum lucidum* Nakai (NC_040920), *Soliva sessilis* Ruiz & Pav. (NC_034851), *Tanacetum cinerariifolium* (Trevir.) Sch.Bip. (MZ984207), *Taraxacum officinale* F.H.Wigg. (NC_030772).




Figure 1. Phylogenetic tree of *Tanacetum cinerariifolium* and related species inferred from maximum likelihood analysis and Bayesian inference based on the complete chloroplast genome sequence. Values above branches indicate bootstrap percentages of maximum likelihood from 1000 replicates, followed by Bayesian posterior probabilities.

Phylogenetic trees were constructed based on complete chloroplast genomes with gene partitioning for common genes using a protocol for standardization of chloroplast sequences described by Turudić et al. (2021). Sequences were aligned using MAFFT v. 7.453 software (Rozewicki et al. 2019), and phylogenetic analyses were performed using maximum likelihood analysis (ML) in RAxML version 8.2.124 (Stamatakis 2014) and Bayesian inference (BI) in MrBayes version 3.2.7a (Ronquist et al. 2012).

The minimum length of the analyzed sequences was 150,136 bp and the maximum length was 152,718 bp. The length of the alignment was 161,421 bp. The number of monomorphic, non-informative, and informative sites in the alignment was 129,382, 8242, and 5954, respectively.

Both maximum likelihood analysis and Bayesian inference yielded identical topologies, with species of the tribe Anthemidae forming a well-supported clade and *T. cinerariifolium* being a sister taxon to species of the genera *Artemisia* and *Chrysanthemum* (Figure 1), consistent with previous studies (Oberprieler et al. 2007; Masuda et al. 2009; Sonboli et al. 2011).

Ethic statement

The collection of plant material was carried out in accordance with guidelines provided by national regulations and permitted by the authority of the Ministry of Environmental Protection and Energy of the Republic of Croatia (UP/1-612-07/17-48/47, 517-07-1-1-1-17-6; 21. April 2017).

Author contributions

AT and ZŠ contributed to the conception and design of the study, analysis, and interpretation of the data. AT wrote the first version of the

manuscript. ZL, MG, JJ, FV, and ZŠ critically reviewed the article regarding its intellectual content. ZL collected biological samples. All authors read, discussed, and approved the final version and all authors agree to be accountable for all aspects of the work.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This research has been funded by the project KK.01.1.1.01.0005 Biodiversity and Molecular Plant Breeding, Center of Excellence for Biodiversity, and Molecular Plant Breeding (CoE CroP-BioDiv), Zagreb, Croatia.

Data availability statement

The genome sequence data that support the findings of this study are openly available in GenBank of NCBI at <https://www.ncbi.nlm.nih.gov> under the accession no. MZ984207. The associated BioProject, SRA, and Bio-Sample numbers are PRJNA799694, SRR17714824, and SAMN25207243, respectively.

References

- Casida JE, Quistad GB. 1995. Pyrethrum flowers: production, chemistry, toxicology, and uses. New York: Oxford University Press.
- Masuda Y, Yukawa T, Kondo K. 2009. Molecular phylogenetic analysis of members of *Chrysanthemum* and its related genera in the tribe Anthemideae, the Asteraceae in East Asia on the basis of the internal transcribed spacer (ITS) region and the external transcribed spacer (ETS) region of nrDNA. *Chromosome Botany*. 4(2):25–36.
- McKain MR, Wilson M. 2018. Fast-plast: rapid de novo assembly and finishing for whole chloroplast genomes. *Fast-Plast v.1.2.8*. Zenodo. <https://github.com/mrmckain/Fast-Plast>.

- Nikolić T. 2020. *Flora croatica*. Alfa d.d. Zagreb Croatia. 2:572.
- Oberprieler C, Himmelreich S, Vogt R. 2007. A new subtribal classification of the tribe Anthemideae (Compositae). *Willdenowia*. 37(1):89.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient bayesian phylogenetic inference and model choice across a large model space. *Syst Biol*. 61(3):539–542.
- Rozewicki J, Li S, Amada KM, Standley DM, Katoh K. 2019. MAFFT-DASH: integrated protein sequence and structural alignment. *Nucleic Acids Res*. 47(W1):W5–W10.
- Sonboli A, Osaloo SK, Vallès J, Oberprieler C. 2011. Systematic status and phylogenetic relationships of the enigmatic *Tanacetum paradoxum* Bornm. (Asteraceae, Anthemideae): evidences from nrDNA ITS, micromorphological, and cytological data. *Plant Syst Evol*. 292(1–2):85–93.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 30(9):1312–1313.
- Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, Greiner S. 2017. GeSeq - versatile and accurate annotation of organellar genomes. *Nucleic Acids Res*. 45(W1):W6–W11.
- Turudić A, Liber Z, Grdiša M, Jakše J, Varga F, Šatović Z. 2021. Towards the well-tempered chloroplast DNA sequences. *Plants*. 10(7):1360.